

Chicken IL-16 VetSet™ ELISA Development Kit



Catalog Number: VS0082C-002
Lot Number: LK0483GK
Quantity: 2 Plate Set
Storage: 2-8°C
Sample Type: Cell Culture Supernatants
Stability: 12 months (from date of receipt)
Country of Origin: USA

Included Components:	Description	Quantity	Component Number	Lot Number
	Chicken IL-16 Coated Plate	2 each	VS0082C-CP	KK0438CJ
	Chicken IL-16 Standard	2 each	VS0082C-ST	JK0419II
	Chicken IL-16 Detection Antibody	2 each	VS0082C-DA	JK0418FJ
	Streptavidin-HRP	1 each	AR0068-001	GK0364HK
	Plate Sealer	6 each	N/A	N/A

Additional Reagents Required:	Reagent	Formulation
	DPBS	0.008M sodium phosphate, 0.002M potassium phosphate, 0.14M sodium chloride, 0.01M potassium chloride, pH 7.4
	Standard and Sample Diluent	Complete cell culture medium used to generate cell culture supernatant samples. It is critical that this medium contain at least 1% carrier protein. If the medium does not contain carrier protein, use Reagent Diluent to dilute the Standard and samples.
	Reagent Diluent	4% BSA in DPBS, 0.2 µm filtered
	Wash Buffer	0.05% Tween®-20 in DPBS
	Substrate	3,3',5,5'-tetramethylbenzidine (TMB) Substrate <i>ELISA Accessory Pack: Catalog # ARO133-002</i>
	Stop Solution	0.18 M Sulfuric Acid <i>ELISA Accessory Pack: Catalog # ARO133-002</i>

Component Preparation:	Component	Preparation
	Chicken IL-16 Standard	Reconstitute Standard in 1 mL Standard and Sample Diluent. Dilute 60 µL of the reconstituted standard in 940 µL of Standard and Sample Diluent. The Standard now has a concentration of 1000 pg/mL . Prepare 1:1 serial dilutions of the Standard by mixing 500 µL Standard with 500 µL Standard and Sample Diluent. Repeat 1:1 serial dilutions until reach a final concentration of 15.6 pg/mL. Use Standard and Sample Diluent as a zero standard.
	Chicken IL-16 Detection Antibody Working Solution	Reconstitute Detection Antibody in 500 µL Reagent Diluent. Dilute the 500 µL of reconstituted Detection Antibody in 11.5 mL Reagent Diluent.
	Streptavidin-HRP Working Solution	Dilute 500 µL of Streptavidin-HRP in 11.5 mL Reagent Diluent.


REAGENTS FOR VETERINARY RESEARCH



Chicken IL-16 VetSet™ ELISA Development Kit

- ELISA Procedure:**
1. Prepare Standard and cell culture supernatant sample dilutions in Standard and Sample Diluent.
 2. Add 100 μ L of Standard or sample to appropriate wells.
Note: Run each Standard or sample in duplicate.
 3. Cover plate with Plate Sealer and incubate at room temperature (20-25°C) for 1 hour.
 4. Wash plate FOUR times with Wash Buffer.
Note: Gently squeeze the long sides of plate frame before washing to ensure all strips remain securely in the frame. Empty plate contents. Use a squirt wash bottle to vigorously fill each well completely with 1X Wash Buffer, then empty plate contents. Repeat procedure three additional times for a total of FOUR washes. Blot plate onto paper towels or other absorbent material.
 5. Add 100 μ L of Detection Antibody Working Solution to each well.
 6. Cover plate with Plate Sealer and incubate at room temperature for 1 hour.
 7. Wash plate FOUR times with Wash Buffer as described in step 4.
 8. Add 100 μ L of Streptavidin-HRP Working Solution to each well.
 9. Cover plate with Plate Sealer and incubate at room temperature for 30 minutes.
 10. Wash plate FOUR times with Wash Buffer as described in step 4.
 11. Add 100 μ L of TMB Substrate Solution to each well.
 12. Develop the plate in the dark at room temperature for 30 minutes.
Note: Do **NOT** cover plate with Plate Sealer.
 13. Stop reaction by adding 100 μ L of Stop Solution to each well.
 14. Measure absorbance on a plate reader at 450 nm.

Typical Standard Curve:



Data represents a typical standard curve generated using the Chicken IL-16 VetSet™ ELISA Development Kit.

A standard curve should be generated with each assay.

Representative Data:

Stimulant	Chicken IL-16 (pg/mL)
Unstimulated	849
Phytohemagglutinin (PHA; 10 μ g/mL)	727
Staphylococcal enterotoxin B (SEB; 5 μ g/mL)	629
Phorbol 12-myristate 13-acetate (PMA; 10 ng/mL) and Ionomycin (500 ng/mL)	1016

PBMCs harvested by ficoll density gradient from an apparently healthy chicken were suspended in RPMI medium containing 10% fetal bovine serum and stimulated as desired. The cell-free supernatants were harvested following three days stimulation and analyzed in the Chicken IL-16 VetSet™ ELISA Development Kit.

Technical Notes: This kit is for the quantitative measurement of Chicken IL-16 in cell culture supernatants. If assaying other sample types, an appropriate Standard and Sample Diluent will need to be developed and validated. Any changes to the ELISA protocol may significantly affect the results generated and will require optimization.

Warranty: Products are warranted by Kingfisher Biotech, Inc. to meet stated product specifications and to conform to label descriptions when used, handled and stored according to instructions. Unless otherwise stated, this warranty is limited to one year from date of sale or expiry date, whichever comes first. Kingfisher Biotech's sole liability for the product is limited to replacement of the product or refund of the purchase price. Kingfisher Biotech products are supplied for research applications. They are not intended for medicinal, diagnostic or therapeutic use. The products may not be resold, modified for resale or used to manufacture commercial products without prior written approval from Kingfisher Biotech, Inc.

VetSet is a trademark of Kingfisher Biotech, Inc.
Tween is a registered trademark of ICI Americas.
© 12/2010

REAGENTS FOR VETERINARY RESEARCH

